

## Does Soil Structure Influence Phosphorus Availability in Andosols? ¿La Estructura del Suelo Influye en la Disponibilidad de Fósforo en Andosoles?

John Clunes<sup>1,4</sup> , Sadi Werner-Auad<sup>1,2</sup> , Daniela Bustos-Korts<sup>2,4</sup> , and  
Felipe Zúñiga-Ugalde<sup>3,4</sup> 

<sup>1</sup> Universidad Austral de Chile, Facultad de Ciencias Agrarias y Alimentarias, Instituto de Ingeniería Agraria y Suelos; (J.C.), (S.W.A.). <sup>2</sup> Instituto de Producción y Sanidad Vegetal; (S.W.A.), (D.B.K.). <sup>3</sup> Facultad de Ciencias Forestales y Recursos Naturales, Instituto de Bosques y Sociedad; (F.Z.U.). <sup>4</sup> Centro de Investigación en Suelos Volcánicos. Independencia 641, Isla Teja. 5091000 Región Los Ríos, Valdivia, Chile; (J.C.), (F.Z.U.).

† Corresponding author: john.clunes@uach.cl

### SUMMARY

Phosphorus (P) availability in the soil is a limiting nutrient for biomass production in agroecosystems developed in Andosols. Nonetheless, it remains unknown how soil structure affects the storage and availability of this nutrient. Thus, this research aimed to evaluate the role of soil structure in phosphorus availability using Olsen P as a chemical indicator of availability. Undisturbed and disturbed soil samples were collected following a grid pattern under naturalized and sowed pastures that reflected two different soil structures (untilled and tilled, respectively). The untilled soil showed higher variability and a higher concentration of Olsen P (5-27 mg kg<sup>-1</sup>) than the tilled soil (14 mg kg<sup>-1</sup>).

Undisturbed samples collected in cylinders showed that Olsen P continues to increase after removing the soil from the cylinder, sieving the sample, and re-extracting Olsen P, resulting in three times the amount of Olsen P found in disturbed samples collected before the experiment. The methodological approach used in this research allowed to highlight the role that soil structure plays in the availability of P over time to improve the efficiency of nutrient utilization.

**Index words:** *allophane, Olsen P, spatial heterogeneity.*

### RESUMEN

La disponibilidad de fósforo (P) en el suelo es un nutriente limitante para la producción de biomasa en agroecosistemas desarrollados en Andosoles. Sin embargo, aún se desconoce si la estructura del suelo afecta el almacenamiento y la disponibilidad de este nutriente. Por lo tanto, esta investigación tuvo como objetivo evaluar el rol de la estructura del suelo en la disponibilidad de fósforo utilizando el P Olsen como indicador. Se tomaron muestras de suelo no disturbadas y disturbadas siguiendo un patrón de cuadrícula bajo praderas naturalizadas y sembradas que reflejaban dos estructuras de suelo diferentes (sin labrar y labrado, respectivamente). El suelo no labrado mostró una mayor variabilidad y concentración de P Olsen (5-27 mg kg<sup>-1</sup>) que el suelo labrado (14 mg kg<sup>-1</sup>).

Las muestras no disturbadas recolectadas en cilindros mostraron que el P Olsen continúa aumentando después de retirar el suelo del cilindro, tamizar la muestra y volver a extraer el P Olsen, lo que da como resultado una cantidad de P Olsen tres veces mayor que la encontrada en las muestras disturbadas recolectadas antes del experimento. El enfoque metodológico utilizado en esta investigación permitió resaltar el papel que juega la estructura del suelo en la disponibilidad de P a lo largo del tiempo para mejorar la eficiencia de la utilización de nutrientes.

**Palabras clave:** *alofán, P Olsen, heterogeneidad espacial.*



#### Cita recomendada:

Clunes, J., Werner-Auad, S., Bustos-Korts, D., & Zúñiga-Ugalde, F. (2025). Does Soil Structure Influence Phosphorus Availability in Andosols? *Terra Latinoamericana*, 43, 1-7. e2047. <https://doi.org/10.28940/terra.v43i.2047>

Received: August 5, 2024.  
Accepted: January 16, 2025.  
Research Note, Volume 43.  
June 2025.

Editor de Sección:  
Dra. Elizabeth Urbina-Sánchez

Editor Técnico:  
M. C. Ayenía Carolina Rosales Nieblas



**Copyright:** © 2025 by the authors.  
Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY NC ND) License (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

## INTRODUCTION

Phosphorus is vital for energy transfer in plants and root development; low amount of available phosphorus may be due to immobilization according to the soil texture (Gen-Jiménez *et al.*, 2025).

Soils derived from volcanic ash are deficient in plant-available phosphorus (P) due to their high retention capacity (> 85%; Valle, Carrasco, Pinochet, Soto, and Mac Donald, 2015; Vásconez, and Pinochet, 2018). It is, therefore, necessary to fertilize these soils with high amounts of P to increase the soil P supply during the growing season (Rodríguez, Pinochet, and Matus, 2001). The soil P availability is often determined through chemical extraction indices that reflect the available P fraction, such as Mehlich 1, Bray 1, and Olsen (Ryan, and Rashid, 2018). These indices have been validated through calibrations between yield and crop P uptake, allowing their use as diagnostic tools for soil fertility (Sandaña, and Pinochet, 2016; Vásconez, and Pinochet, 2018). However, the soil has a heterogeneous distribution of available P content, i.e., there will be soil spots with a high P content and other areas with a low P content, which contrasts with the assumption that the soil is homogeneous before any extraction (Pinochet, 1995<sup>1</sup>). P is an immobile nutrient in the soil, i.e., phosphates have low solubility, short distance diffusion, and depend on the adsorption sites. It is assumed that these adsorption sites are not dependent on soil structure. Therefore, we hypothesized that soil structure (i.e., a network porous system) is a key factor in soil functions (Bronick, and Lal, 2005), providing substantial amounts of available P, and therefore, P availability determined through Olsen P overestimates available P over time. This research aimed to evaluate the role of soil structure on phosphorus availability in pastures under two levels of structuring (tilled and untilled).

## MATERIALS AND METHODS

Soil samples were collected at a pasture at the Estación Experimental Agropecuaria Austral (EEAA) of the Universidad Austral de Chile (39° 46' S, 73° 13' W, 12 meters of altitude) located in the city of Valdivia, Chile. The mean annual temperature there is between 11-12 °C and the mean annual precipitation is between 1800-2400 mm (González-Reyes, and Muñoz, 2013; Dörner *et al.*, 2022).

Southern Chilean soils derived from volcanic ash have developed in mesic temperature and udic moisture regimes are classified as Duric Hapludand (CIREN, 2003) or Silandic Andosol (IUSS Working Group WRB, 2022), are characterized by a highly reactive non-crystalline clay fraction (i.e., allophane and imogolite) (Clunes, and Pinochet, 2021). These soils have values of extractable aluminum in ammonium acetate of > 800 mg kg<sup>-1</sup> (Clunes, Dörner, and Pinochet, 2021), NaF pH values of ≥ 9.4 (Valle *et al.*, 2015), pH in water 5.4-5.8 (Zúñiga *et al.*, 2023), high soil organic matter content (> 12%; Matus, Rumpel, Neculman, Panichini, and Mora, 2014; Bravo *et al.*, 2020) and low bulk density < 0.9 Mg m<sup>-3</sup> (Dörner *et al.*, 2022).

### Soil Sampling and Design

To evaluate the effect of pasture management on soil structure and P availability, disturbed soil samples were collected with an auger, and undisturbed soil samples were collected in steel cylinders (h = 5.60 cm and Ø = 7.15 cm) at a depth of 0-20 cm (depth commonly used to evaluate soil fertility; Rodríguez *et al.*, 2001 and which represents the genetic Ap horizon; Bravo *et al.*, 2020). Samples were collected from two pasture conditions: i) naturalized-degraded pasture > 10 years or untilled (T1) and ii) sown pasture < 1 year or tilled (T2). To evaluate the spatial dependency of P availability, the sampling in T1 and T2 was conducted following a grid pattern (20 m × 20 m) using the center of the plots as a reference.

### Laboratory Analyses

For the disturbed soil samples, the Olsen P methodology (Olsen P, classic) was used to determine soil P availability (Sadzawka *et al.*, 2006). However, for the undisturbed soil samples, modifications to the classic methodology were made to adjust the analysis (Olsen P, cylinder). Briefly, the cylinders with soil were placed in Buchner funnels and connected to a vessel that received an extractant solution of NaHCO<sub>3</sub> (0.5 M - pH 8.5). For each sample, 4 L of NaHCO<sub>3</sub> was applied, maintaining the soil:solution ratio (1:2.5) of the classical methodology (Sadzawka *et al.*, 2006) to avoid variations in the chemical equilibrium during the extraction process. The ratio was adjusted considering the bulk density of the soil (0.75 Mg m<sup>-3</sup> according to Dörner *et al.*, 2022). The solution was applied to the soil using drippers, which allowed homogeneous percolation inside the cylinder with soil. Once all the solution had been received, it was homogenized and filtered. Then, 5 mL of the filtrate was collected in a glass container and 20 mL of the color development reagent was added. Finally, it was left to stand for 60 min for the colorimetric reading of molybdenum blue using a spectrophotometer (Sadzawka *et al.*, 2006). The activated charcoal application and shaking process were not carried out.

<sup>1</sup> Pinochet, D., (1995). *The residual effect of applications of phosphate fertilizer measured by the Olsen method*. Doctoral dissertation, Thesis of Doctor of Philosophy, The University of Reading, United Kingdom.

## Olsen P Statistical Analysis

The Olsen P extraction method (classic and cylinder) and the soil conditions (tilled and untilled) were assessed with a two-way ANOVA with interaction using the 'lm' procedure in R Studio (R Core Team, 2020). Means and standard errors were estimated with the package 'emmeans' (Lenth, 2023).

Olsen P spatial dependency was modeled for each combination of tillage levels and P extraction method with a 2-dimensional P-splines linear mixed model, as proposed by Boer (2023), and implemented in the R-Package LMMsolver as indicated in Equation (1).

$$p_{ij} = \mu + x_i + y_j + s(x)_i + s(y)_j + \epsilon_{ij} \quad (1)$$


Where  $p_{ij}$  is the observation at the  $i^{\text{th}}$  position on the row coordinates and the  $j^{\text{th}}$  position on the columns.  $x_i$  and  $y_j$  are the row and column positions, which are fitted as linear covariates.  $s(x)_i$  is the smooth (p-spline) component along the rows and  $s(y)_j$  is the smooth (p-spline) component along the columns with 20 segments in both directions.  $\epsilon_{ij}$  is a homogeneous residual.

Results were visualized as; i) the difference between the predicted surfaces and the mean of the observed Olsen P at each combination of extraction method and soil condition (expressed as a percentage of the mean), and ii) the difference between the predicted surface for the classic and the cylinder method.

## RESULTS AND DISCUSSION

### Comparison of Olsen P Classic and Olsen P Cylinder Methods

Olsen P was measured with the classic method and in a soil cylinder. There was a strong effect of the soil processing method ( $P < 0.001$ , the mean of the classical method was  $14.2 \text{ mg kg}^{-1}$ , the mean of the cylinder method was  $9.4 \text{ mg kg}^{-1}$ , and the s.e.m -  $0.6 \text{ mg kg}^{-1}$ ). In contrast, the main effect of the tillage and the soil by tillage interactions were non-significant. Furthermore, the difference between the classical method and the cylinder method was of similar magnitude, regardless of the soil and the Olsen P level, as shown by the points below the 1:1 line in Figure 1 (except for two samples that had much higher Olsen P values at the classic method and that corresponded to samples obtained in untilled soil). When removing both extreme points that were observed in the classic method, both methods were positive and linearly associated ( $P = 0.048$ , Figure 1). However, the slope in the equation was small ( $y = 3.91 + 0.48x$ , Figure 1), suggesting that the Olsen P effectively available (cylinder method) might be lower than the Olsen P obtained by the classic method. The higher values for the classic method suggest that soil sample homogenization destroys the soil aggregates and releases the Olsen P stored, making it more detectable during the quantification procedure. The Olsen P determination in cylinders showed that while the concentration of available P in the tilled samples decreased (between  $4$  and  $17 \text{ mg kg}^{-1}$ ), the variability in the untilled increased (Figure 2C). Soil structure, as a parameter of soil function, greatly affects nutrient cycling, thus influencing the nutrients available to roots (Vogel *et al.*, 2018; Clunes *et al.*, 2021). Soil structure indicators, such as aggregate stability, water movement, and pore connectivity, allow for the partial assessment of this function because nutrient storage and recycling also depend on soil chemical properties (Rabot, Wiesmeier, Schlü, and Vogel, 2018). Therefore, P could be stored in the soil both chemically and physically because P is indirectly involved in soil aggregation. While it is true that we do not provide measurements of soil vegetation cover, below-ground biomass production, colonization of arbuscular mycorrhizal fungi, and the formation of phosphate bonding agents, we recognize these properties as valuable elements that help to understand the role of physical protection in the P availability in the soil (Bronick, and Lal, 2005; Borie *et al.*, 2019). It is essential to relate the effect of soil structure on P availability in pastures because an average estimate from a representative number of samples (homogenizing the soil condition and disrupting the soil aggregates) results in an overestimation of available P and therefore an over-fertilization, which leads to an unbalanced and inefficient nutrient use. This effect is supported by the differences in Olsen P concentration, which was obtained in extreme conditions of the experiment ( $14 \text{ mg kg}^{-1}$  in tilled-classic vs.  $8 \text{ mg kg}^{-1}$  untilled-cylinder), as shown in Table 1.



**Figure 1. Association between the Olsen P determined by the classic method (Olsen P, classic) in disturbed soil samples (Tilled) and the Olsen P determined in undisturbed soil samples (Olsen P, cylinder; Untilled).** The black line represents the 1:1 relationship.

### Spatial variation of soil Olsen P


Although there was variation in the Olsen P for the four combinations of soil Olsen P, the spatial dependence was very low. This is reflected in the low effective dimensions for the linear and spline spatial components, and in the predicted surfaces (Figure 2). In most cases, the variation was low, with ranges within the  $\pm 20\%$  variation, relative to the mean of the observed Olsen P, leading to standard deviations of 7.8 % (tilled, classic, Figure 2A), 7.2 % (tilled, cylinder, Figure 2B) and 7.5 % (untilled, cylinder, Figure 2D). The exception was one point in the untilled, classical, which had 28.4 mg kg<sup>-1</sup> Olsen P, leading to a standard deviation of 42.2% (expressed as a percentage of difference compared to the mean of the observations, Figure 2C).

It has been reported that factors such as reduced pasture growth due to drought or lack of irrigation in the summer, water accumulation in areas of the pasture during winter, poor grazing frequency and less palatable species can result in reduced nutrient removal from the pasture and hence greater spatial variability of P in a permanent pasture (Cotching, Taylor, and Corkrey, 2020). In addition, P accumulation in the upper centimeters of soil in the degraded permanent pasture may also be due to the prolonged period during which this soil was undisturbed (Nze-Memiaghe, Cambouris, Ziadi, Karam, and Perron, 2021), which for this study was over 10 years (Descalzi, López, Kemp, Dörner, and Ordóñez, 2020). In general, a sampling depth of 0 to 20 cm is recommended for soil fertility tests, leading to a mixing of soil layers; in the case of untilled permanent pastures, the spatial variability of P presents points with higher P concentrations (Toor *et al.*, 2020). In this context, the decision to collect 0 to 20 cm disturbed soil samples aims to define whether the soil has the Olsen P concentration required for adequate pasture nutrition, which, for Andosols, should be around 20 mg kg<sup>-1</sup> (Vistoso, Iraira, and Sandaña, 2021).

**Table 1. Multiple comparison among Olsen P means for tilled and untilled pastures.**

Pasture condition	Method	Olsen P mean
Tilled	Classic	14.17 A**
	Cylinder	10.64 AB
Untilled	Classic	11.96 AB
	Cylinder	8.12 B**

\*\* Indicate significance differences between means ( $P = 0.0032$ ).



**Figure 2. P-spline predicted surfaces for Olsen P for A) Tilled soil, classical method, B) Tilled soil, cylinder method, C) Untilled, classical method, and D) Untilled, cylinder method.** Color scale is proportional to the percentage of change at each grid position, relative to the mean of the predicted surface. The mean and the range of the predicted surface are indicated below the title for each combination of soil and method. The standard deviation is expressed as the percentage of change in relation to each mean.

This critical value is due to the formation of complexes between Al- or Fe- and colloidal material (organic carbon, allophane, and imogolite) that permit the retention of large quantities of soil organic P (Redel *et al.*, 2016). Werner *et al.* (2017) reported that in soil aggregates from P-rich areas, P was co-located with aluminum, iron oxides, and hydroxides, while in soil aggregates from P-depleted areas, the phosphorous was bonded to the soil organic carbon. This would be related to P accretion areas or small “pockets” through a process of re-sorption in soil aggregates (Pinochet, 1995). This hypothesis has not yet been probed, but it is an interesting approach that supports the idea of physical protection of P proposed in this preliminary study. Therefore, the amount of soil P available for plants should be above the critical level at which the crop does not respond to the application of P fertilizer, which ensures a high crop yield without causing severe risks of contaminating the agroecosystem (Díaz, and Torrent, 2016).

Pastures on volcanic soils in southern Chile have high organic P storage (848-1065 mg kg<sup>-1</sup>), and the availability of this nutrient is mainly regulated by the formation of amorphous Al-Po complexes (Redel *et al.*, 2016). Phosphate fertilizer applications increase the concentration of available P in the soil solution, which allows for a rapid diffusion of P to the root system and, thus, an increased P uptake by plants (Vistoso *et al.*, 2021). However, underestimating the initial P content in the soil using chemical extraction methods that do not consider the soil structure in which the root system grows and from which it absorbs P, results in an inefficient and far from rational application of phosphate fertilizers.

## CONCLUSIONS

Through the approach presented, this preliminary report seeks to spotlight the importance of soil structure on nutrient availability, especially for P, an immobile and highly retention nutrient in volcanic soils.

Soil structure plays a relevant role in the capacity to deliver available P over space and time. The above can be particularly relevant in agroecosystems that promote soil conservation, including zero tillage, natural pastures such as steppes, and sown areas when aggregate formation begins.

We are conscious that the limitations of this preliminary experiment should be considered for future research in the area, such as the relationship between soil P availability rate and plant P uptake, site climatic conditions, and soil P supply during the growing season.

## ETHICS STATEMENT

Not applicable.

## CONSENT FOR PUBLICATION

Not applicable.

## AVAILABILITY OF SUPPORTING DATA

The datasets used and analyzed during the current study are available from the corresponding author upon reasonable request.

## COMPETING INTERESTS

The authors declare that they have no competing interests.

## FINANCING

Initiation FONDECYT grant 11221038 provided by ANID, Chile.

## AUTHORS' CONTRIBUTIONS

Conceptualization, J.C. and F.Z.; methodology, J.C., F.Z. and S.W.; validation, F.Z. and D.B.; formal analysis, F.Z. and D.B.; investigation, J.C., S.W., and F.Z.; data curation, F.Z. and D.B.; writing-original draft preparation, J.C.; writing-review and editing, J.C., S.W., F.Z., and D.B.; visualization, J.C., S.W., F.Z., and D.B.; supervision, J.C. and F.Z.

## ACKNOWLEDGMENTS

The first author thanks the FONDECYT grant 11221038 (Initiation FONDECYT) provided by ANID for funding this research project.

## REFERENCES

- Boer, M. P. (2023). Tensor product P-splines using a sparse mixed model formulation. *Statistical Modelling*, 23(5-6), 465-479. <https://doi.org/10.1177/1471082X231178591>
- Borie, F., Aguilera, P., Castillo, C., Valentine, A., Seguel, A., Barea, J. M., & Cornejo, P. (2019) Revisiting the Nature of Phosphorus Pools in Chilean Volcanic Soils as a Basis for Arbuscular Mycorrhizal Management in Plant P Acquisition. *Journal of Soil Science and Plant Nutrition*, 19, 390-401. <https://doi.org/10.1007/s42729-019-00041-y>
- Bravo, S., González-Chang, M., Dec, D., Valle, S., Wendroth, O., Zúñiga, F., & Dörner, J. (2020). Using wavelet analyses to identify temporal coherence in soil physical properties in a volcanic ash-derived soil. *Agricultural and Forest Meteorology*, 285, 107909. <https://doi.org/10.1016/j.agrformet.2020.107909>
- Bronick, C. J., & Lal, R. (2005). Soil structure and management: a review. *Geoderma*, 124(1-2), 3-22. <https://doi.org/10.1016/j.geoderma.2004.03.005>
- CIREN (Centro de Información de Recursos Naturales). (2003). *Estudio agrológico - X Región: descripciones de suelos, materiales y símbolos*. Santiago, Chile: CIREN. ISBN: 956-7153-48-5

- Clunes, J., & Pinochet, D. (2021). Leucine retention by the clay-sized mineral fraction. An indicator of C storage. *Agro Sur*, 48, 37-46. <https://doi.org/10.4206/agrosur.2020.v48n3-05>
- Clunes, J., Dörner, J., & Pinochet, D. (2021). How does the functionality of the pore system affects inorganic nitrogen storage in volcanic ash soils? *Soil and Tillage Research*, 205, 104802. <https://doi.org/10.1016/j.still.2020.104802>
- Cotching, W. E., Taylor, L., & Corkrey, S. R. (2020). Spatial variation of soil nutrients in dairy pasture paddocks. *New Zealand Journal of Agricultural Research*, 63(4), 492-504. <https://doi.org/10.1080/00288233.2019.1576212>
- Descalzi, C. A., López, I. F., Kemp, P. D., Dörner, J., & Ordóñez, I. (2020). Pasture restoration improvement methods for temperate degraded pastures and consequences of the climatic seasonality on soil-pasture complex. *Journal of Agronomy and Crop Science*, 206(1), 130-147. <https://doi.org/10.1111/jac.12368>
- Díaz, I., & Torrent, J. (2016). Changes in Olsen P in relation to P balance in contrasting agricultural soils. *Pedosphere*, 26(5), 636-642. [https://doi.org/10.1016/S1002-0160\(15\)60072-8](https://doi.org/10.1016/S1002-0160(15)60072-8)
- Dörner, J., Bravo, S., Stoorvogel, M., Dec, D., Valle, S., Clunes, J., ... & Zúñiga, F. (2022). Short-term effects of compaction on soil mechanical properties and pore functions of an Andisol. *Soil and Tillage Research*, 221, 105396. <https://doi.org/10.1016/j.still.2022.105396>
- Gen-Jiménez, A., Sarmiento-Megchum, E. F., Maldonado-Gómez, J. C., Manzano-Gómez, L. A., Solís-Zebadúa, S., Rincón-Molina, C. I., & Rincón-Rosales, R. (2025). Bacterias Rizosféricas y Endófitas con Potencial PGPB Aisladas del Cultivo de Maíz (*Zea mays* L.) en Chiapas, México. *Terra Latinoamericana*, 43, 1-14. <https://doi.org/10.28940/terra.v43i.2002>
- González-Reyes, Á., & Muñoz, A. A. (2013). Cambios en la precipitación de la ciudad de Valdivia (Chile) durante los últimos 150 años. *Bosque (Valdivia)*, 34(2), 200-213. <http://dx.doi.org/10.4067/S0717-92002013000200008>
- IUSS Working Group WRB. (2022). *International soil classification system for naming soils and creating legends for soil maps* (4<sup>th</sup> ed.). Vienna, Austria: International Union of Soil Sciences. ISBN 979-8-9862451-1-9
- Lenth, R. V. (2023). *emmeans: Estimated Marginal Means, aka Least-Squares Means*. Vienna, Austria: R Foundation for Statistical Computing.
- Matus, F., Rumpel, C., Neculman, R., Panichini, M., & Mora, M. L. (2014). Soil carbon storage and stabilisation in andic soils: A review. *Catena*, 120, 102-110. <https://doi.org/10.1016/j.catena.2014.04.008>
- Nze-Memighe, J. D., Cambouris, A. N., Ziadi, N., Karam, A., & Perron, I. (2021). Spatial variability of soil phosphorus indices under two contrasting grassland fields in eastern Canada. *Agronomy* 11(1), 24. <https://doi.org/10.3390/agronomy11010024>
- Rabot, E., Wiesmeier, M., Schlüter, S., & Vogel, H. J. (2018). Soil structure as an indicator of soil functions: A review. *Geoderma*, 314, 122-137. <https://doi.org/10.1016/j.geoderma.2017.11.009>
- Redel, Y., Cartes, P., Demanet, R., Velásquez, G., Poblete-Grant, P., Bol, R., & Mora, M. L. (2016). Assessment of phosphorus status influenced by Al and Fe compounds in volcanic grassland soils. *Journal of Soil Science and Plant Nutrition*, 16(2), 1-17. <http://dx.doi.org/10.4067/S0718-95162016005000041>
- Rodríguez, J., Pinochet, D., & Matus, F. (2001). Fertilización de los cultivos. Santiago, Chile: LOM ediciones. ISBN: 978-956-288-880-6
- R Core Team (2020). *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing.
- Ryan, J., & Rashid, A. (2005). Phosphorus. In R. Lal (Ed.), *Encyclopedia of Soil Science* (2<sup>nd</sup> ed.) (pp. 1275-1279). Boca Raton, FL, USA: CRC Press.
- Sandaña, P., & Pinochet, D. (2016). Phosphorus acquisition of wheat, pea and narrow-leafed lupin under different P supplies. *Journal of Soil Science and Plant Nutrition*, 16(2), 537-549. <http://dx.doi.org/10.4067/S0718-95162016005000044>
- Sadzawka, A., Carrasco, M. A., Grez, R., Mora, M. L., Flores, H., & Neaman, A. (2006). Recommended methods of analysis for soils in Chile. *Commission Standards and Accreditation. Chilean Society of Soil Science*. Santiago, Chile: Instituto de Investigaciones Agropecuarias.
- Toor, G. S., Yang, Y. Y., Morris, M., Schwartz, P., Darwish, Y., Gaylord, G., & Webb, K. (2020). Phosphorus pools in soils under rotational and continuous grazed pastures. *Agrosystems, Geosciences & Environment*, 3(1), 1-12. <https://doi.org/10.1002/agg2.20103>
- Valle, S. R., Carrasco, J., Pinochet, D., Soto, P., & Mac Donald, R. (2015). Spatial distribution assessment of extractable Al, (NaF) pH and phosphate retention as tests to differentiate among volcanic soils. *Catena*, 127, 17-25. <https://doi.org/10.1016/j.catena.2014.12.011>
- Vásconez, G., & Pinochet, D. (2018). Residual value of the phosphate added to ecuadorian and chilean soils with different phosphorus retention capacity. *Journal of Soil Science and Plant Nutrition*, 18(1), 60-72. <http://dx.doi.org/10.4067/S0718-95162018005000301>
- Vistoso, E., Iraira, S., & Sandaña, P. (2021). Phosphorus use efficiency in permanent pastures in Andisols. *Journal of Soil Science and Plant Nutrition*, 21, 2587-2599. <https://doi.org/10.1007/s42729-021-00526-9>
- Vogel, H. J., Bartke, S., Daedlow, K., Helming, K., Kögel-Knabner, I., Lang, B., ... & Wollschläger, U. (2018). A systemic approach for modeling soil functions. *Soil*, 4(1), 83-92. <https://doi.org/10.5194/soil-4-83-2018>
- Werner, F., Mueller, C. W., Thieme, J., Gianoncelli, A., Rivard, C., Höschel, C., & Prietzel, J. (2017). Micro-scale heterogeneity of soil phosphorus depends on soil substrate and depth. *Scientific Reports*, 7(3203), 1-9. <https://doi.org/10.1038/s41598-017-03537-8>
- Zúñiga, F., Leiva, C., Rivano, F., San Martín, M. & Toledo, P. (2023). Propiedades de suelos distribuidos en un transecto longitudinal del Sur de Chile. *Agro Sur*, 51(3), 1-8. <https://doi.org/10.4206/agrosur.2023.v51n3-01>